


University of Belgrade
Institute of Physics Belgrade
Kopaonik, March 13-16, 2022



Book of Abstracts
15th Photonics Workshop
(Conference)



15th Photonics Workshop (2022)

Book of abstracts

Kopaonik, Serbia, March 13-16, 2022

Publisher, 2022:

Institute of Physics Belgrade

Pregrevica 118

11080 Belgrade, Serbia

Editors:

Dragan Lukić, Marina Lekić, Zoran Grujić

ISBN 978-86-82441-55-7

Printed by:

NEW IMAGE d.o.o.

Tošin Bunar 185, Belgrade

Number of copies: 55

CIP - Каталогизација у публикацији - Народна библиотека Србије, Београд

535(048)

681.7(048)

66.017/.018(048)

PHOTONICS Workshop (15; 2022; Kopaonik)

Book of Abstracts / 15th Photonics Workshop, (Conference), Kopaonik,
March 13-16, 2022; [editors Dragan Lukić, Marina Lekić, Zoran Grujić]. -

Belgrade: Institute of Physics, 2022 (Belgrade: New image). - 72 str.:

ilustr.; 25 cm

Tiraž 55. - Registar.

ISBN 978-86-82441-55-7

a) Оптика - Апстракти b) Оптиелектроника - Апстракти c) Технички
материјали - Апстракти

COBISS.SR-ID 60055049

Anisotropy of dielectric permittivity in optical gain modelling of MIR quantum cascade lasers

Nikola Vuković¹, Jelena Radovanović¹, Vitomir Milanović¹

(1) *University of Belgrade, School of Electrical Engineering, Bulevar kralja Aleksandra 73, 11120 Serbia*

Contact: N. Vuković (nvukovic@etf.bg.ac.rs)

Abstract. An enhanced numerical model for calculation of population densities and optical gain in quantum cascade laser (QCL) [1] that takes into account anisotropy of dielectric permittivity [2] of the layered quantum-well based structure is presented. Our results indicate that the layering induced anisotropy leads to significant changes in electron-longitudinal (LO) phonon scattering rates [3] on account of the existence of additional resonant frequencies for LO phonons, each altering the total scattering rate between the relevant subbands. Namely, instead of taking LO phonon frequency as a constant [4], we have included dependence on phonon wavevector in the vicinity of the LO phonon modes: one dependence originates from the material system that is used and the others on the transition energies of the structure's active region. Subsequently, carrier distribution between the subbands and the degree of population inversion in the mid-infrared QCL active region are substantially changed in comparison to the predictions of an isotropic permittivity model widely used in the literature [5]. Numerical calculations presented in this contribution are based on solving the system of rate-equations iteratively and then evaluating carrier densities, scattering rates, and optical gain.

REFERENCES

- [1] J. Faist, Quantum Cascade Lasers. OUP Oxford (2013).
- [2] N. Vuković et al., Opt. Quantum Electron. 47, 883–891 (2015).
- [3] M. A. Stroschio and M. Dutta, Phonons in Nanostructures, Cambridge University Press, (2001).
- [4] C. Jirauschek and T. Kubis, Appl. Phys. Rev. 1, p. 011307, (2014).
- [5] P. Harrison, A. Valavanis, Quantum Wells, Wires and Dots, John Wiley & Sons, Ltd, Chichester, UK (2016).

Acknowledgments

The authors acknowledge support from Ministry of Education, Science and Technological Development (Republic of Serbia), "Multi-Scale Modeling of Terahertz Quantum Cascade Laser Active Regions", Multilateral scientific and technological cooperation in the Danube region 2020-2021, and "DEMETRA: Development of high-performance mid-IR / THz quantum cascade lasers for advanced applications", Science Fund of the Republic of Serbia, Serbian Science and Diaspora Collaboration Programme: Knowledge Exchange Vouchers.